

AD-A146 134


HIGH TEMPERATURE GAS ENERGY TRANSFER(U) WASHINGTON UNIV 1/1
SEATTLE DEPT OF CHEMISTRY B S RABINOVITCH 07 SEP 84
N00014-75-C-0690

UNCLASSIFIED

F/G 7/4

NL

END
DATE
FILMED
10-84
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963-A

AD-A146 134

ANNUAL SUMMARY REPORT, 1 August 1983 - 31 July 1984

High Temperature Gas Energy Transfer

B. S. Rabinovitch, Principal Investigator

Department of Chemistry BG-10
University of Washington
Seattle, Washington 98195

OFFICE OF NAVAL RESEARCH

Contract N00014-75-C-0690

Project NR 092-549

7 September 1984

DTIC
ELECTE
OCT 3 1984
S B

DTIC FILE COPY

This document has been approved for public release; its distribution is unlimited. Reproduction in whole or in part is permitted for any purpose of the United States Government. This research was sponsored by the Office of Naval Research.

84 . 09 28 021

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
	AD-A146134	
4. TITLE (and Subtitle)	5. TYPE OF REPORT & PERIOD COVERED	
High Temperature Gas Energy Transfer	Annual Summary 1 Aug 1983 - 31 Jul 1984	
	6. PERFORMING ORG. REPORT NUMBER	
7. AUTHOR(s)	8. CONTRACT OR GRANT NUMBER(s)	
B. S. Rabinovitch	N00014-75-C-0690 NR 092 549	
9. PERFORMING ORGANIZATION NAME AND ADDRESS	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS	
Department of Chemistry BG-10 University of Washington Seattle WA 98195		
11. CONTROLLING OFFICE NAME AND ADDRESS	12. REPORT DATE	
Office of Naval Research, Code 432 Department of the Navy 800 N Quincy. Arlington VA 22217	7 Sept 1984	
	13. NUMBER OF PAGES	
	16	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	15. SECURITY CLASS. (of this report)	
	15a. DECLASSIFICATION/DOWNGRADING SCHEDULE	
16. DISTRIBUTION STATEMENT (of this Report)		
This document has been approved for public release; its distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)		
Energy Transfer Thermal System Gases Transients Metal Surface Variable Encounter Method High Temperature Vibrational Energy Liquid Surface Wall Activation		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)		
Study of the transfer of vibrational energy between molecules and surfaces was continued at high levels of vibrational excitation and high surface temperature. Past work is summarized. Reports TR28 and TR29 were issued in this period. Metal and liquid surfaces have been examined. Results are outlined.		

DD FORM 1473
1 JAN 73EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102 LF 014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

I. Summary of Progress and Future Plans

Work on vibrational energy transfer at a surface by "energetic" species has been completed. Work on the last of these, azomethane, was reported in this period. Other energetic species previously studied included azomethane and methyl nitrite. We confirm the results of our earlier work, namely:

Apart from a propensity for more complex reaction mechanisms and catalytic behavior, the energy transfer properties of energetic materials at a surface follow the behavior expected from the intermolecular potential and not that of the reaction potential. More stable molecules can be used to delineate the energy transfer properties expected for energetic materials.

Work on vibrational energy transfer by gas molecules at a metal surface has been extended from the case of a solid (gold) surface to liquid (tin) surface. We confirm and extend our earlier conclusion drawn from the gold surface, namely:

Metal surfaces are stronger vibrational energy transfer media than pyrex or silica surfaces. They are all ideal strong collision agents at low temperatures (< 400-600 K).

To this we add a new finding, namely:

No discontinuity in collisional gas/surface energy transfer efficiency results from change of phase at the metal melting point.

Finally, theoretical treatment of a Boltzmann strong collision model has led to generalizations on energy transfer efficiency as a function of both molecule size, ambient temperature and hot molecule energy level.

Future plans include completion of work in progress on gallium liquid metal surfaces, application of time-of-flight studies to other molecule(s) (methyl cyclobutene), and study of ionic salt surfaces as vibrational energy transfer agents.

PREVIOUS PAGE
IS BLANK



Dist	Avail and/or Special
A-1	

II. Complete Listing of Technical Reports.

1. Technical Report TR01, 7 January 1976.

Vibrational Energy Transfer in Reaction Systems at Elevated Temperatures by B. S. Rabinovitch, D. G. Keil, J. F. Burkhalter and G. B. Skinner.

In this report a theoretical analysis was given of shock tube data for high temperature gas reactions. The data show that collisional energy transfer efficiency appears to decrease at high temperatures.

2. Technical Report TR02, 20 April 1976

Temperature Dependence of the Arrhenius Activation Energy. High Temperature Limit, J. F. Burkhalter and B. S. Rabinovitch.

In this report a conventional approximation to the high temperature vibrational partition function is shown to be defective, and the usual high temperature expression for the Arrhenius activation is in error. The variation of the activation energy for unimolecular reactions as a function of temperature is calculated for various representative systems.

3. Technical Report TR03, 15 October 1976

Intermolecular Vibrational Energy Transfer in Thermal Unimolecular Systems, D. C. Tardy and B. S. Rabinovitch.

This report gives a comprehensive and detailed survey and analysis of energy transfer in thermal gaseous systems. The present status of the field is delineated and future areas of work indicated.

4. Technical Report TR04, 15 April 1977

On the Use of Exact Vibrational State Counting Methods in RRKM Rate Calculations, S. E. Stein and B. S. Rabinovitch.

This report describes an algorithm for accurate state sum and density calculations and refutes an error in the literature.

5. Technical Report TR05, 15 August 1977

Vibrational Translational Energy Transfer in Atom-Polyatomic Molecule Collisions in Thermal Reaction Systems by I. Oref and B. S. Rabinovitch.

In this report a simple theoretical model describing energy transfer probabilities is given. Conservation of angular momentum was imposed. The importance of completeness and detailed balance conditions and correction of an earlier model in the literature are illustrated. The model is applied to experimental data.

6. Technical Report TR06, 1 September 1977

Problems of Diffusion in a Low Pressure Gas Stream as Related to the Polanyi Diffusion Flame (Cloud) Method by D. G. Keil, J. F. Burkhalter and B. S. Rabinovitch.

A theoretical and experimental study was made of flow and diffusion of a reactant in a low pressure gas stream. Connection is made with earlier theoretical analysis.

7. Technical Report TR07, 10 October 1978

Do Highly Excited Reactive Polyatomic Molecules Behave Ergodically?
I. Oref and B. S. Rabinovitch.

The literature is surveyed and analyzed. A variety of excitation techniques including crossed molecular beam, laser, chemical activation, photochemical and thermal are considered. The answer to the title question is "yes." The relationship to the present work on intermolecular transfer is the following: multiphoton laser experiments frequently involve collisional heating of the gas. In order to understand the results, one must have a good basis for interpretation of data. In this survey, some suggested models for energy relaxation by molecular collisions are criticized and data reinterpreted in light of results obtained in this laboratory.

8. Technical Report TR08. 1 November 1978

Vibrational Energy Transfer in Thermal Unimolecular Systems by the Diffusion Cloud Method. Cyclopropane. E. Kamaratos, J. D. Burkhalter, D. G. Keil and B. S. Rabinovitch.

A study of vibrational energy transfer by the Diffusion Cloud Method has been made at temperatures from 975 K to 1175 K in the cyclopropane isomerization system. H_2 , He, N_2 and CO_2 were studied as inert bath gases. Their relative efficiencies increase in that order. Values of $\langle \Delta E \rangle$ vary from 150 cm^{-1} to 1100 cm^{-1} at 975 K. It was shown that these values decline, and the collisional efficiencies, β_c , decrease markedly with rise of temperature to 1175 K. This is only the second measurement of such a phenomenon.

9. Technical Report TR09. 15 March 1979

Collisional Relaxation of Non-Equilibrium Vibrational Energy Distributions in a Thermal Unimolecular System. Surface Collisions. D. F. Kelley, B. D. Barton, L. Zalotai, and B. S. Rabinovitch.

A novel, simple technique, the Variable Encounter Method, is described for obtaining information on energy transfer efficiency between a gas and a solid in the transient region, and as a function of temperature. The method is as powerful as it is simple. Results for cyclopropane are described. Decrease

in collisional efficiency $\langle \Delta E \rangle$ with rise of temperature is confirmed, and the form of the transfer probability matrix \tilde{P} is discovered. This method and these results are considered to be some of the most innovative and important that have ever been made in this field.

10. Technical Report TR10. 15 June 1979

Transients in Vibrational Relaxation of Polyatomic Molecules at High Temperatures by VEM. M. C. Flowers, D. F. Kelley, F. C. Wolters, and B. S. Rabinovitch.

Extension of the VEM technique to cyclobutane is described. An exponential form of \tilde{P} is required to fit the data. The value of $\langle \Delta E \rangle$ declines with rise of temperature.

11. Technical Report TR11; 15 September 1979

Collisional Relaxation of Transient Vibrational Energy Distributions in a Thermal Unimolecular System. The Variable Encounter Method. D. F. Kelley, L. Zalotai and B. S. Rabinovitch.

The Variable Encounter Method permits the study of the transient in the relaxation of an initial vibrationally cold ensemble of molecules in a vibrationally hot distribution by a known and variable number of successive collisions with a hot wall. The system studied was the isomerization of 1,1-cyclopropane- d_2 with a fused quartz wall temperature of 800 K to 1175 K, and average number of collisions from 2.3 to 22.3. The detailed theory of the experiment was presented, together with the method of data deconvolution. Various modified gaussian and exponential models of energy transfer were found to give agreement with the data. The average down-step size was found to decline from $\leq 3500 \text{ cm}^{-1}$ at the lowest temperature to $\sim 2500 \text{ cm}^{-1}$ at the highest on the basis of a gaussian model, but heterogeneous energy transfer is more efficient than homogeneous gas-gas collisional transfer. A mathematical analysis of the relation between mean first-passage times and incubation times is given. Incubation times increase from ~ 7 to ~ 12 collisions with increasing temperature. Transient population distributions and the sequential reaction probabilities as a function of collision number are calculated.

12. Technical Report TR12. 10 October 1979

Vibrational Energy Transfer in a Diffusion-Flow Cyclopropane- d_2 System. J. F. Burkhalter, E. Kamaratos, and B. S. Rabinovitch.

In this study, absolute rate constants for isomerization of cyclopropane- d_2 and high temperature homogeneous energy transfer were determined in a diffusion-flow system. The relative rates of the competitive isotopic isomerization channels

were measured for two bath gases, N_2 and He, at two temperatures, 973 K and 1073 K. Values of the average energy down-jump size $\langle \Delta E \rangle$ were computed from both the absolute rates and the isotopic relative rates by suitable modelling by a stochastic calculation. The results confirm earlier diffusion cloud measurements.

13. Technical Report TR13, 10 October 1979

Transients in the Vibrational Excitation of Cyclobutane Decomposition Using the Variable Encounter Method. M. C. Flowers, F. C. Wolters, B. D. Barton and B. S. Rabinovitch.

The probability of reaction of cyclobutane molecules in a fixed-time interval after experiencing a known number of collisions with a hot surface at temperatures between 749 K and 1126 K was determined using the Variable Encounter Method. Calculations utilizing exponential or gaussian models for energy transfer enabled the average amounts of energy transferred for deactivating collisions, $\langle \Delta E' \rangle$, to be estimated. The exponential model fits the experimental data best and, using this model, $\langle \Delta E' \rangle$ is 2430 cm^{-1} at 748 K and decreases to 1470 cm^{-1} at 1123 K. Surface collisions are more efficient than binary gas collisions; but cyclobutane is a less efficient partner than cyclopropane- d_2 (TR No. 11). The results follow from a quasi-statistical model of energy accommodation.

14. Technical Report TR14, 1 December 1979

A Crucial Demonstration of Strong Collisional Behavior of Vibrational Energy for Gas-Surface Collisions. M. C. Flowers, F. C. Wolters, D. F. Kelley and B. S. Rabinovitch.

Application of the VEM technique was made to the thermal cyclobutene decomposition system. Vibrational energy transfer in the cyclobutene - seasoned quartz surface system was studied. Collisional efficiency again declines with rise of temperature, but the occurrence of strong collisions was demonstrated in crucial manner at temperatures below 450 K. The wall appears to behave like a theoretical strong collider below 450 K. This is a very novel finding.

15. Technical Report TR15, 31 March 1980

Gas-Surface Vibrational Energy Transfer in the Transient Region of a Low-Pressure Unimolecular Reaction. B. D. Barton, D. F. Kelley and B. S. Rabinovitch.

The Variable Encounter Method, for the study of gas-wall vibrational energy transfer in the transient region of a unimolecular reaction was applied to the

isomerization of cyclopropane to propylene. Temperatures in the range 900 K - 1125 K were employed. The average probability of reaction per collision, $\bar{P}_c(m)$, was deduced from the data and compared with a theoretical stochastic calculation based on both gaussian and exponential models for the energy transfer probabilities. The former model is more appropriate. The efficiency of a seasoned quartz wall was greater than gas-gas collisions of substrate and the efficiency declined with increase of temperature. The steady state is closely approached (90%) in a comparatively small number of gas-wall collisions - 10-20, approximately. The efficiency of transfer by cyclopropane is comparable with that by cyclopropane- d_2 (TR 11).

16. Technical Report TR16, 1 August 1980

Collisional Relaxation of Vibrational Energy Transients in the Methyl Cyclopropane System. D. F. Kelley, T. Kasai and B. S. Rabinovitch.

The variable encounter method has been used to study vibrational energy transients in the isomerization of methylcyclopropane to various butenes. This molecule, which has a hindered methyl rotation mode, has the highest vibrational energy level density of any molecule - cyclopropane, cyclobutene, cyclobutane - that has been studied so far in testing the variation of energy transfer efficiency with molecular structure. This system was studied with reactor surface temperatures of 800 to 1130 K and average numbers of collisions per encounter with the reactor of 5.6 and 20.0. An exponential model of energy transfer was found to give the best fit to the data with the average down step energy $\langle \Delta E' \rangle$ decreasing from 1860 to 1415 cm^{-1} with increase of temperature over the range studied. Incubation times increased from 14 to 19 collisions with increase in temperature, and these times, together with values of the conventional relative collision efficiency β and values of $\langle \Delta E' \rangle$, are compared with those of other molecules studied by VEM. The calculated transient population distributions and the associated sequential reaction probabilities are also displayed. The energy transfer efficiency from the hot molecules in down-collisions is lowest in this system and hence the weakest model - exponential - gives a good fit. By contrast, the efficiency of up-transitions, i.e., energy transfer into the cold molecule measured in this work is higher than for other molecules. This is a consequence of the increased

eigenstate density and the important conclusion is reached that this verifies the suitability of a statistical model of energy accommodation to describe vibrational energy transfer involving high energy polyatomic molecules at these temperatures.

17. Technical Report TR17, 15 September 1980

A Reinvestigation of Transients in the Cyclopropane System by the Variable Encounter Method. M. C. Flowers, F. C. Wolters, D. F. Kelley and B. S. Rabinovitch.

The isomerization of cyclopropane to propene was reinvestigated under conditions identical to those used in a previously reported VEM study on cyclobutane. The simultaneous reaction of cyclobutane was also studied in the largest reactor. Some details of the method are amplified; the distribution functions for numbers of collisions in each reactor are displayed as are the contributions to reaction $R(n)$ as a function of the number of consecutive collision n . The present data are in essential agreement with those obtained by Kelley et al [TR 11], although the average size of an internal energy down transition ($\langle \Delta E' \rangle$) for cyclopropane molecules colliding with a hot surface is a little lower than previously estimated. The increasing efficiency of the surface in deactivating energized molecules as the surface temperature decreases is confirmed $\langle \Delta E' \rangle = 2550 \text{ cm}^{-1}$ at 900 K, on a gaussian model for energy transfer, and $\langle \Delta E' \rangle = 2000 \text{ cm}^{-1}$ at $\sim 1100 \text{ K}$. The surface acts as a somewhat stronger collider for cyclopropane than for cyclobutane and is also a more efficient collider for cyclopropane than are gas-gas cyclopropane collisions. This accords with the results of TR16 and leads to the important conclusion that heterogeneous collision is more efficient than homogeneous collision, in general, for the deactivation of excited molecules.

18. Technical Report TR18, 15 October 1980

Transients in Thermal Isomerization of Cyclobutene by the Variable Encounter. Vibrational Energy Transfer and Relaxation at Lower Temperatures. F. C. Wolters, M. C. Flowers and B. S. Rabinovitch.

Vibrational energy transfer and relaxation in the thermal isomerization of cyclobutene to 1,3-butadiene was studied over the temperature range 429 K - 778 K using the Variable Encounter Method. Reactors characterized by various mean number of collisions, m , per encounter of a substrate molecule with the reactor ($m = 2.6, 5.9, 8.5$ and 27.2) were used. The

significant observation was made that strong collider behavior in collisions of substrate with a seasoned fused quartz wall is reached at lower temperatures (below 450 K), as evidenced by the coincidence of values of the probability of reaction per collision, $\bar{P}_c(m)$, obtained in VEM reactors having different m values. The average energy transferred to the wall per deactivating collision, $\langle \Delta E' \rangle$, increased strongly as temperature decreased, from $\langle \Delta E' \rangle = 2150 \text{ cm}^{-1}$ at 764 K to $> 6000 \text{ cm}^{-1}$ at 460 K. Various forms of models for the probability of energy transfer in down transitions were applied to data. The weak form — exponential — cannot apply, in contrast to the behavior that applies at higher temperatures for looser molecules (higher vibrational eigenstate densities) such as methyl cyclopropane (TR16).

19. Technical Report TR19, 15 March 1981.

Energy Transfer at High Temperatures in the Cyclopropane System.
Temperature Dependence of $\langle \Delta E' \rangle$. T. Kasai, D. F. Kelley and B. S. Rabinovitch.

The study of collisional energy transfer between cyclopropane molecules and a seasoned silica wall by the Variable Encounter Method was extended over a range of more than 500° upward in temperature to 1325 K. The value of $\langle \Delta E' \rangle$, the average energy of a down-transition upon collision of the hot substrate molecule with the wall to decrease monotonically. The data provide the most accurate and well-documented display for a polyatomic molecule of the temperature trend of $\langle \Delta E' \rangle$ presently available. The rate of decrease in $\langle \Delta E' \rangle$ seems to be more rapid at lower temperatures. The data have been explained by the theory of Lin and Rabinovitch on a quasi-statistical accommodation model.

The behavior at still higher temperatures is considered to be not well known and must be considered as in quite unsatisfactory condition. It is of extreme practical and theoretical importance to detect the possible transition of the observed behavior from an attractive potential-dominated accommodation phenomenon to a Landau-Teller type repulsive potential-dominated impulsive phenomenon.

20. Technical Report TR20, March 15, 1981

Single-Collision Gas-Surface Vibrational Energy Transfer in a Reacting System. D. F. Kelley, T. Kasai and B. S. Rabinovitch.

Gas-surface vibrational energy accommodation in a reactive system has been studied under single collision conditions for the first time for polyatomic molecules. The reaction system was the isomerization of cyclobutene to 1,3-butadiene. Both seasoned pyrex and silica surfaces were used over the temperature ranges 400-775 K and 500-975 K, respectively. Strong collider behavior was observed below ~ 425 K. The vibrational energy accommodation coefficient was found to decrease with rise of temperature from ~ 1.0 to 0.2, while the relative collisional efficiency β_1 declined from ~ 1.0 to 0.008. A stochastic calculation was used to fit the data. This technique provides a more sensitive test between the various model forms for energy transfer than has been possible heretofor. It is clearly shown that a Gaussian or particularly a Boltzmann weighted exponential form provide the best fit.

21. Technical Report TR21, August 1, 1981

Pyrolysis of 1-Iodopropane by the Variable Encounter Method. F. C. Wolters, K.-J. Chao and B. S. Rabinovitch

The pyrolysis of 1-iodopropane has been studied by the Variable Encounter Method (VEM) at temperatures from 625 K to 840 K. Deactivating wall collisions are found to be stronger for this molecule than for the hydrocarbons previously studied in this temperature range. This is attributed to the increased strength of the attractive molecule-surface interaction due to the dipole moment and enhanced polarizability of the substrate molecule.

22. Technical Report TR22, October 20, 1981

Isotope Effect in Gas-Surface Vibrational Energy Transfer. Cyclopropane and Cyclopropane- d_6 Isomerization by the VEM Technique. W. Yuan, R. Tosa, K.-J. Chao, and B. S. Rabinovitch

The study of the effect of varying molecular structure upon the efficiency of vibrational energy transfer between initially cold molecules and a hot surface was extended to the reaction pair, cyclopropane/cyclopropane- d_6 . The latter member is somewhat less efficient in energy loss. This was interpreted in terms of molecular properties, namely, the latter's greater vibrational eigenstate.

density. Conversely, the probability of energy up-transitions is enhanced. In thermal low pressure unimolecular reactions, this corresponds to the inverse statistical weight secondary isotope effect. Comparison was made with other molecules.

23. Technical Report TR23, November 25, 1981

Single Collision Gas-Surface Vibrational Energy Transfer in Reactive Systems. Variation of Initial Energy Distribution. R. Arakawa, D. F. Kelley, and B. S. Rabinovitch

The initial vibrational energy distribution of molecules that collide with a hot surface, which is an important parameter of the interaction, has been studied under single collision conditions. These experiments provide a more sensitive test of the relative suitability of various analytical forms for the collisional transition probability matrix, \tilde{P} . The reaction system is the isomerization of cyclobutene to 1,3 butadiene. A seasoned fused quartz surface was used over the temperature range $T_r = 600 \text{ K} - 900 \text{ K}$. Variation of the initial vibrational energy population vector of cyclobutene molecules was made by change of their initial temperature T_c in the range $273 \text{ K} - 620 \text{ K}$. Gaussian or Boltzmann Exponential forms of \tilde{P} prove to be the most suitable to fit the data.

24. Technical Report TR24, January 15, 1982

Vibrational Energy Transfer and Pyrolysis of Nitromethane by the Variable Encounter Method. W. Yuan, B. S. Rabinovitch and R. Tosa

Nitromethane is of considerable interest because of its large dipole moment and importance in a variety of chemical systems. The pyrolysis of nitromethane has been studied by the Variable Encounter Method (VEM) at temperatures from 816 K to 1092 K with two reactors of differing geometry having fused silica surfaces. The probability of reaction per collision with the reactor surface was measured. The down energy transition jump size, $\langle \Delta E' \rangle$, was determined. It decreased with increasing wall temperature. A comparison is made of $\langle \Delta E' \rangle$ with previous results reported to date for other substrate molecules. Indeed, nitromethane is one of the more efficient energy transfer agents, as might be expected from the strength of the attractive interaction. However, in addition to the size (vibrational eigenstate density) and the polarity of the molecules, the nature of the hot surface seems also to play a role.

25. Technical Report TR25, June 1, 1982

The Isomerization of 1-Methyl Cyclobutene by the Single Collision Variable Encounter Method. R. Arakawa and B. S. Rabinovitch

The reaction probability, P_c , for 1-methylcyclobutene molecules energized at a hot silica surface was measured under single collision conditions over the temperature range $T_r = 480 \text{ K} - 800 \text{ K}$. The initial thermal vibrational energy population of the cyclobutene was varied from $T_c = 273 \text{ K} - 570 \text{ K}$. The reaction was isomerization to isoprene. These experiments provide a test of the relative suitability of various analytical forms for the collisional transition probability matrix \tilde{P} . Calculations with a Gaussian form provide the best overall fit to the data. The average amount of energy ($\langle \Delta E \rangle_{E_0}$) transferred from the hot molecules in a vibrational down transition from the reaction threshold energy level, E_0 , declined from 7220 cm^{-1} to 3890 cm^{-1} with increase in surface temperature from 600 K to 800 K. The experimental collisional efficiency, β_1 , declined from 0.39 to 0.035 over the combination temperature range $T_r, T_c = 600, 500$ to $800, 293$. Strong collider behavior was observed for T_r less than 450 K.

The Following were submitted in the present report period:

26. Technical Report TR26, December 1, 1982

Vibrational Energy Transfer at a Gold Surface in Reacting Systems. Cyclobutene and Nitromethane. Wei Yuan and B. S. Rabinovitch

Single collision excitation probabilities were measured for cyclobutene and nitromethane on polycrystalline gold plane and wire surfaces. Transport above the reaction thresholds for isomerization and decomposition, respectively, for the two substrates was used as the criterion of vibrational energy transfer. Several different seasoning and processing procedures of the surfaces were tested. For cyclobutene, a defect from strong collider transition probabilities appeared only above 550 K - 600 K, which signifies that the energy transfer efficiency at a treated gold surface is greater than that at a seasoned silica surface. The vibrational energy transfer efficiency declined above 600 K and fell abruptly to a quasi-constant value in the range 900 K - 1100 K. Above 1100 K the differences between various conditioned surfaces diminished progressively. Unlike cyclobutene, and earlier studies of nitromethane on silica surfaces where reproducible non-catalytic behavior could be attained, nitromethane on gold showed apparent catalytic phenomena which the various conditioning treatments failed to repress.

27. Technical Report TR27, 31 July 1983.

Study of Vibrational Energy Transfer at a Surface by a Time-of-Flight Method. Wei Yuan and B.S. Rabinovitch.

A single collision, time-of-flight extension of the VEM method for the study of molecule-surface vibrational energy transfer is introduced. This refined technique helps election between possible alternative trial analytic forms of the collisional transition probability function. A gaussian form is preferred over a Boltzmann-exponential form for cyclobutene isomerization to 1,3-butadiene energized by collisions at a seasoned silica surface at 800K. The study illustrates the difference between conventional accommodation coefficient measurements, which examine average energy changes, and the present studies which explore the energy transport at high levels corresponding to chemical reaction thresholds. It confirms the hopes for refined exploration of the transition probability function expressed in section IV D of the Annual Report issued in August 1982.

28. Technical Report TR28, 1 October 1983.

Average Vibrational Energy Transfer During a Single Collision of Excited Molecules with Heat Bath Molecules, I. Dref and B.S. Rabinovitch.

A collisional energy transfer probability of the form $B(E+\Delta E) \int_{-\infty}^{\infty} B'(E') B'(E'-\Delta E) \delta E'$ is assumed where $B(E)$ is the Boltzmann distribution and ΔE is the incremental energy transferred and which can display negative as well as positive values. Single collisions between vibrationally excited substrate molecules with heat bath molecules are considered. The dependence of the average energy per collision transferred up, down and overall on the initial energy content, on the temperature and on the size of the bath and substrate molecules is calculated and compared with experimental data in the literature.

29. Technical Report TR29, 10 January 1984.

Vibrational Energy Transfer at a Surface in Reacting Systems. Decomposition of Azomethane. W. Nilsson, E. Linkaityte-Weiss and B.S. Rabinovitch.

Single collision activation of the "energetic" species azomethane at a hot seasoned fused silica surface was studied. Excitation above the critical threshold ($E_0 \approx 50 \text{ kcal mole}^{-1}$) was monitored by subsequent homogeneous decomposition. The collisional efficiency was studied over the range 700K to 875K. Energy transfer efficiency is discussed and compared with values

for other substrate molecules of comparable size and polarity. The reaction thermodynamics do not affect the energy transfer properties which depend on the inter- rather than intra-molecular potential. The data emphasize the uncertainty that still prevails with respect to the values of the Arrhenius parameters for this system.

III. Listing of Publications.

1. Vibrational Energy Transfer in Reaction Systems at Elevated Temperatures, Proceed. Tenth Internat'l. Shock Tube Sympos. ed. G. Kamimoto, Kyoto, 1976.
2. Intermolecular Vibrational Energy Transfer in Thermal Unimolecular Systems, D.C. Tardy and B.S. Rabinovitch, *Chem. Revs.* 77, 369 (1977).
3. On the Use of Exact Vibrational State Counting Methods in RRKM Rate Calculations, S.E. Stein and B.S. Rabinovitch, *Chem. Phys. Lett.* 49, 183 (1977).
4. Vibrational Translational Energy Transfer in Atom-Polyatomic Molecule Collisions in Thermal Reaction Systems, I. Dref and B.S. Rabinovitch, *Chem. Phys.* 26, 385 (1977).
5. Problems of Diffusion in a Low Pressure Gas Stream as Related to the Polany Diffusion Flame (Cloud) Method, D.G. Keil, J.F. Burkhalter and B.S. Rabinovitch, *J. Phys. Chem.* 83, 984 (1979).
6. Vibrational Energy Transfer in Thermal Unimolecular Systems by the diffusion Cloud Method. Cyclopropane, E. Kamaratos, J.F. Burkhalter, D.G. Keil and B.S. Rabinovitch, *J. Phys. Chem.* 83, 984 (1979).
7. Do Highly Excited Reactive Polyatomic Molecules Behave Ergodically? I. Dref and B.S. Rabinovitch, *Accts. Chem. Res.* 12, 166 (1979).
8. Collisional Relaxation of Non-equilibrium Vibrational Energy Distributions in a Thermal System. Surface Collisions, D.F. Kelley, B.D. Barton, L. Zalotai and B.S. Rabinovitch, *J. Chem. Phys.* 71, 538 (1979).
9. Transients in Vibrational Relaxation of Polyatomic Molecules at High Temperatures by Variable Encounter Method, M.C. Flowers, B.D. Barton, F.C. Wolters and B.S. Rabinovitch, *Amer. Soc. Mass. Spectrom. Proceed.*, ed. A.E. Harrison, 27, 661 (1979).
10. Collisional Relaxation of Transient Vibrational Energy Distributions in a Thermal Unimolecular System. The Variable Encounter Method, D.F. Kelley, L. Zalotai, and B.S. Rabinovitch, *Chem. Phys.* 46, 379 (1980).
11. Vibrational Energy Transfer in a Diffusion-Flow Cyclopropane-d₂ System, J.F. Burkhalter, E. Kamaratos and B.S. Rabinovitch *J. Phys. Chem.* 84, 476 (1980).
12. A Crucial Demonstration of Strong Collisional Behavior of Vibrational Energy for Gas-Surface Collision. M.C. Flowers, F.C. Wolters, D.F. Kelley and B.S. Rabinovitch, *Chem. Phys. Lett.* 69, 543 (1980).
13. Transients in the Vibrational Excitation of Cyclobutane Decomposition Using the Variable Encounter Method, M.C. Flowers, F.C. Wolters, B.D. Barton and B.S. Rabinovitch, *Chem. Phys.* 47, 189 (1980).
14. Gas-Surface Vibrational Energy Transfer in the Transient Region of a Low-Pressure Unimolecular Reaction. B.D. Barton, D.F. Kelley, and B.S. Rabinovitch, *J. Phys. Chem.* 84, 1299 (1980).

15. Collisional Relaxation of Vibrational Energy Transients in the Methylcyclopropane System. A Variable Encounter Method Study, D.F. Kelley, T. Kasai and B.S. Rabinovitch, 73, 5611 (1980).
16. Transients in Thermal Isomerization of Cyclobutene by VEM. Vibrational Energy Transfer and Relaxation at Lower Temperatures, *J. Phys. Chem.* 85, 589 (1981).
17. Surface-Gas Energy Transfer in the Cyclopropane/Cyclobutane Isomerization Systems by the Variable Encounter Method, M.C. Flowers, F.C. Wolters, D.F. Kelley and B.S. Rabinovitch, *J. Phys. Chem.* 85, 849 (1981).
18. Single-Collision Gas-Surface Vibrational Energy Transfer in a Reacting System, D.F. Kelley, T. Kasai and B.S. Rabinovitch, *J. Phys. Chem.* 85, 1100 (1981).
19. Energy Transfer at High Temperatures in the Cyclopropane System. Temperature Dependence of $\langle \Delta E' \rangle$, T. Kasai, D.F. Kelley and B.S. Rabinovitch, *Chem. Phys. Lett.* 81, 126 (1981).
20. Pyrolysis of 1-Iodopropane by the Variable Encounter Method, F.C. Wolters, K.-J. Chao and B.S. Rabinovitch, *Internat'l J. Chem. Kin.* 13, 1289 (1981).
21. Isotope Effect in Gas-Surface Vibrational Energy Transfer. Cyclopropane and Cyclopropane- d_6 Isomerization by the TEM Technique. W. Yuan, R. Tosa, K.-J. Chao and B.S. Rabinovitch, *Chem. Phys. Lett.* 85, 27 (1982).
22. Single Collision Gas-Surface Vibrational Energy Transfer in Reactive Systems. Variation of Initial Energy Distribution. R. Arakawa, D.F. Kelley and B.S. Rabinovitch, *J. Chem. Phys.* 76, 2384 (1982).
23. Vibrational Energy Transfer and Pyrolysis of Nitromethane by the Variable Encounter Method. W. Yuan, B.S. Rabinovitch and R. Tosa, *J. Phys. Chem.* 86, 2796 (1982).
24. The Isomerization of 1-Methyl Cyclobutene by Single Collision Activation at a Surface. Variation of Initial Energy, R. Arakawa and B.S. Rabinovitch, *J. Phys. Chem.* 86, 4772 (1982).
25. Vibrational Energy Transfer at a Gold Surface in a Reacting System. Cyclobutene and Nitromethane, W. Yuan and B.S. Rabinovitch, *J. Phys. Chem.* 81, 2167 (1983).
26. Study of Vibrational Energy Transfer at a Surface by a Time-of-Flight Method, W. Yuan and B.S. Rabinovitch, *J. Chem. Phys.* 80, 1687 (1984).
27. Average Vibrational Energy Transfer During Single Collision of Excited Molecules with Heat Bath Molecules. I. Drefand and B.S. Rabinovitch, *J. Chem. Soc. Faraday Trans., I* 80, 769 (1984).

IV. Other Work Completed or in Progress and Future Work.

In addition to the accomplishment summarized in Technical Reports TR28 and TR29, the following work has been completed or is in progress.

A. Liquid Metals and Other Surfaces.

As was pointed out in an earlier summary, it is of practical and theoretical interest to investigate energy transfer at other surfaces. Very little work reported in the literature on molecule/liquid surface energy transfer. In 1937, Alty (Sci. Progr. 31, 436 (1937)) reported that the accommodation coefficient α is unity for water(g)/water(l), Hg(g)/Hg(l), CCl₄(g)/CCl₄(l), and C₂H₅OH(g)/C₂H₅OH(l). Since then, in one study in 1967 (unpublished), Thomas has reported that α for He(g)/K(l) is less than unity. The above work was carried out near room temperature and, to our knowledge, high temperature energy transfer at liquid metal surfaces has not been studied. We have completed such work with tin which is now being issued as technical report TR30. Single collision vibrational energy transfer for a canonical ensemble (T=300K) of cyclobutene molecules at a liquid tin surface was studied at temperatures from 500K to 775K. Transport above the reaction threshold for isomerization to butadiene ($E_0 = 32.4$ kcal mole⁻¹) was used as the criterion for efficiency of vibrational energy accommodation. The surface was found to function as a strong collider below 550K. Experiments both above and below the freezing point revealed no discontinuity in the vibrational accommodation efficiency. This is discussed in terms of the structure of liquid and solid tin.

We now have additional studies underway with gallium liquid surfaces and can extend the studies to salt surfaces.

B. Time-of-Flight Studies.

In TR27, 31 July 1983, we reported on the development and application of a time-of-flight method to the elucidation of the functional form of the collision transition probability matrix P in the study of vibrational energy transfer by cyclobutene gas molecules at a seasoned silica surface. We are now extending this work to the more favorable case of methyl cyclobutene which is experimentally advantageous for two reasons: a) this molecule provides a more continuous vibrational eigenstate density distribution at low energies; and b) it has a lower rate of decomposition at given energy which provides more refined delineation of functional form in our apparatus, as discussed in TR27.

INIT

DISTRIBUTION LIST

7 September 84

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. L.V. Schmidt Assistant Secretary of the Navy (R,E, and S) Room 5E 731 Pentagon Washington, D.C. 20350	1	Dr. F. Roberto Code AFRPL MKPA Edwards AFB, CA 93523	1
Dr. A.L. Slafkosky Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1	Dr. L.H. Caveny Air Force Office of Scientific Research Directorate of Aerospace Sciences Bolling Air Force Base Washington, D.C. 20332	1
Dr. Richard S. Miller Office of Naval Research Code 432 Arlington, VA 22217	10	Mr. Donald L. Ball Air Force Office of Scientific Research Directorate of Chemical Sciences Bolling Air Force Base Washington, D.C. 20332	1
Mr. David Siegel Office of Naval Research Code 260 Arlington, VA 22217	1	Dr. John S. Wilkes, Jr. FJSRL/NC USAF Academy, CO 80840	1
Dr. R.J. Marcus Office of Naval Research Western Office 1030 East Green Street Pasadena, CA 91106	1	Dr. R.L. Lou Aerojet Strategic Propulsion Co. P.O. Box 15699C Sacramento, CA 95813	1
Dr. Larry Peebles Office of Naval Research East Central Regional Office 666 Summer Street, Bldg. 114-D Boston, MA 02210	1	Dr. V.J. Keenan Anal-Syn Lab Inc. P.O. Box 547 Paoli, PA 19301	1
Dr. Phillip A. Miller Office of Naval Research Naval Station, Treasure Island Bldg. 7, Rm. 81 San Francisco, CA 94130	1	Dr. Philip Howe Army Ballistic Research Labs ARRADCOM Code DRDAR-BLT Aberdeen Proving Ground, MD 21005	1
Mr. Otto K. Heiney AFATL - DLDL Eglin AFB, FL 32542	1	Mr. L.A. Watermeier Army Ballistic Research Labs ARRADCOM Code DRDAR-BLI Aberdeen Proving Ground, MD 21005	1
Mr. R. Geisler ATTN: MKP/MS24 AFRPL Edwards AFB, CA 93523	1	Dr. W.W. Wharton Attn: DRSMI-RKL Commander U.S. Army Missile Command Redstone Arsenal, AL 35898	1

PRECEDING PAGE BLANK-NOT FILLED

INIT

DISTRIBUTION LIST

	<u>No. Copies</u>		<u>No. Copies</u>
Mr. J. Murrin Naval Sea Systems Command Code 62R2 Washington, D.C. 20362	1	Dr. A. Nielsen Naval Weapons Center Code 385 China Lake, CA 93555	1
Dr. P.J. Pastine Naval Surface Weapons Center Code R04 White Oak Silver Spring, MD 20910	1	Dr. R. Reed, Jr. Naval Weapons Center Code 388 China Lake, CA 93555	1
Mr. L. Roslund Naval Surface Weapons Center Code R122 White Oak Silver Spring, MD 20910	1	Dr. L. Smith Naval Weapons Center Code 3205 China Lake, CA 93555	1
Mr. M. Stosz Naval Surface Weapons Center Code R121 White Oak Silver Spring, MD 20910	1	Dr. B. Doua Naval Weapons Support Center Code 5042 Crane, IN 47522	1
Dr. E. Zimmet Naval Surface Weapons Center Code R13 White Oak Silver Spring, MD 20910	1	Dr. A. Faulstich Chief of Naval Technology MAT Code 0716 Washington, D.C. 20360	1
Dr. D.R. Derr Naval Weapons Center Code 388 China Lake, CA 93555	1	LCDR J. Walker Chief of Naval Material Office of Naval Technology MAT, Code 0712 Washington, D.C. 20360	1
Mr. Lee N. Gilbert Naval Weapons Center Code 3205 China Lake, CA 93555	1	Mr. Joe McCartney Naval Ocean Systems Center San Diego, CA 92152	1
Dr. E. Martin Naval Weapons Center Code 3858 China Lake, CA 93555	1	Dr. S. Yamamoto Marine Sciences Division Naval Ocean Systems Center San Diego, CA 91232	1
Mr. R. McCarten Naval Weapons Center Code 3272 China Lake, CA 93555	1	Dr. G. Bosmajian Applied Chemistry Division Naval Ship Research & Development Center Annapolis, MD 21401	1
		Dr. H. Shuey Rohm and Haas Company Huntsville, AL 35801	1

INIT

DISTRIBUTION LIST

	<u>No. Copies</u>		<u>No. Copies</u>
Mr. R. Brown Naval Air Systems Command Code 330 Washington, D.C. 20361	1	Dr. J. Schnur Naval Research Lab. Code 6510 Washington, D.C. 20375	1
Dr. H. Rosenwasser Naval Air Systems Command AIR-310C Washington, D.C. 20360	1	Mr. R. Beauregard Naval Sea Systems Command SEA 64E Washington, D.C. 20362	1
Mr. B. Sobers Naval Air Systems Command Code 03P25 Washington, D.C. 20360	1	Mr. G. Edwards Naval Sea Systems Command Code 62R3 Washington, D.C. 20362	1
Dr. L.R. Rothstein Assistant Director Naval Explosives Dev. Engineering Dept. Naval Weapons Station Yorktown, VA 23691	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, PA 19112	1
Dr. Lionel Dickinson Naval Explosive Ordnance Disposal Tech. Center Code D Indian Head, MD 20640	1	Dr. H.G. Adolph Naval Surface Weapons Center Code R11 White Oak Silver Spring, MD 20910	1
Mr. C.L. Adams Naval Ordnance Station Code PM4 Indian Head, MD 20640	1	Dr. T.D. Austin Naval Surface Weapons Center Code R16 Indian Head, MD 20640	1
Mr. S. Mitchell Naval Ordnance Station Code 5253 Indian Head, MD 20640	1	Dr. T. Hall Code R-11 Naval Surface Weapons Center White Oak Laboratory Silver Spring, MD 20910	1
Dr. William Tolles Dean of Research Naval Postgraduate School Monterey, CA 93940	1	Mr. G.L. Mackenzie Naval Surface Weapons Center Code R101 Indian Head, MD 20640	1
Naval Research Lab. Code 6100 Washington, D.C. 20375	1	Dr. K.F. Mueller Naval Surface Weapons Center Code R11 White Oak Silver Spring, MD 20910	1

INIT

6/81

DISTRIBUTION LIST

	<u>No. Copies</u>		<u>No. Copies</u>
Dr. R.G. Rhoades Commander Army Missile Command DRSMI-R Redstone Arsenal, AL 35898	1	Dr. E.H. Debutts Hercules Inc. Baccus Works P.O. Box 98 Magna, UT 84044	1
Dr. W.D. Stephens Atlantic Research Corp. Pine Ridge Plant 7511 Wellington Rd. Gainesville, VA 22065	1	Dr. James H. Thacher Hercules Inc. Magna Baccus Works P.O. Box 98 Magna, UT 84044	1
Dr. A.W. Barrows Ballistic Research Laboratory USA ARRADCOM DRDAR-BLP Aberdeen Proving Ground, MD 21005	1	Mr. Theodore M. Gilliland Johns Hopkins University APL Chemical Propulsion Info. Agency Johns Hopkins Road Laurel, MD 20810	1
Dr. C.M. Frey Chemical Systems Division P.O. Box 358 Sunnyvale, CA 94086	1	Dr. R. McGuire Lawrence Livermore Laboratory University of California Code L-324 Livermore, CA 94550	1
Professor F. Rodriguez Cornell University School of Chemical Engineering Olin Hall Ithaca, NY 14853	1	Dr. Jack Linsk Lockheed Missiles & Space Co. P.O. Box 504 Code Org. 83-10, Bldg. 154 Sunnyvale, CA 94088	1
Defense Technical Information Center DTIC-DDA-2 Cameron Station Alexandria, VA 22314	12	Dr. B.G. Craig Los Alamos National Lab P.O. Box 1663 NSP/DOD, MS-245 Los Alamos, NM 87545	1
		Dr. R.L. Rabie WX-2, MS-952 Los Alamos National Lab. P.O. Box 1663 Los Alamos, NM 37545	1
Dr. Ronald L. Simmons Hercules Inc. Eglin AFATL/DLDL Eglin AFB, FL 32542	1	Dr. R. Rogers Los Alamos Scientific Lab. WX-2 P.O. Box 1663 Los Alamos, NM 87545	1

6/81

DISTRIBUTION LIST

<u>No. Copies</u>	<u>No. Copies</u>
Dr. J.F. Kincaid 1 Strategic Systems Project Office Department of the Navy Room 901 Washington, D.C. 20376	Dr. C.W. Vriesen 1 Thiokol Elkton Division P.O. Box 241 Elkton, MD 21921
Strategic Systems Project Office 1 Propulsion Unit Code SP2731 Department of the Navy Washington, D.C. 20376	Dr. J.C. Hinshaw 1 Thiokol Wasatch Division P.O. Box 524 Brigham City, UT 83402
Mr. E.L. Throckmorton 1 Strategic Systems Project Office Department of the Navy Room 1048 Washington, D.C. 20376	U.S. Army Research Office 1 Chemical & Biological Sciences Division P.O. Box 12211 Research Triangle Park, NC 27709
Dr. D.A. Flanigan 1 Thiokol Huntsville Division Huntsville, AL 35807	Dr. R.F. Walker 1 USA ARRADCOM DRDAR-LCE Dover, NJ 07801
Mr. G.F. Mangum 1 Thiokol Corporation Huntsville Division Huntsville, AL 35807	Dr. T. Sinden 1 Munitions Directorate Propellants and Explosives Defense Equipment Staff British Embassy 3100 Massachusetts Ave. Washington, D.C. 20008
Mr. E.S. Sutton 1 Thiokol Corporation Elkton Division P.O. Box 241 Elkton, MD 21921	Mr. J.M. Frankle 1 Army Ballistic Research Labs ARRADCOM Code DRDAR-BLI Aberdeen Proving Ground, MD 21005
Dr. G. Thompson 1 Thiokol Wasatch Division MS 240 P.O. Box 524 Brigham City, UT 84302	Dr. Ingo W. May 1 Army Ballistic Research Lab ARRADCOM Code DRDAR-BLI Aberdeen Proving Ground, MD 21005
Dr. T.F. Davidson 1 Technical Director Thiokol Corporation Government Systems Group P.O. Box 9258 Ogden, UT 84409	

INIT

6/81

DISTRIBUTION LIST

	<u>No. Copies</u>		
E. J. Palm Commander Army Missile Command DRSMI-RK Redstone Arsenal, AL 35898	1	Dr. Kenneth O. Hartman Hercules Aerospace Division Hercules Incorporated Allegany Ballistics Lab P.O. Box 210 Cumberland, MD 21502	1
Dr. Merrill K. King Atlantic Research Corp. 5390 Cherokee Avenue Alexandria, VA 22314	1	Dr. Joyce J. Kaufman The Johns Hopkins University Department of Chemistry Baltimore, MD 21218	1
Dr. R.J. Bartlett Batelle Columbus Laboratories 505 King Avenue Columbus, OH 43201	1	Dr. John K. Dienes T-3, MS-216 Los Alamos National Lab P.O. Box 1663 Los Alamos, NM 87544	1
Dr. P. Rentzepis Bell Laboratories Murray Hill, NJ 07971	1	Dr. H.P. Marshall Dept. 52-35, Bldg. 204.2 Lockheed Missile & Space Co. 3251 Hanover Street Palo Alto, CA 94304	1
Professor Y.T. Lee Department of Chemistry University of California Berkeley, CA 94720	1	Professor John Deutsch MIT Department of Chemistry Cambridge, MA 02139	1
Professor M. Nicol Department of Chemistry 405 Hilgard Avenue University of California Los Angeles, CA 90024	1	Professor Barry Kunz College of Sciences & Arts Department of Physics Michigan Technological Univ. Houghton, MI 49931	1
Professor S.S. Penner University of California Energy Center Mail Code B-010 La Jolla, CA 92093	1	Dr. R. Bernecker Code R13 Naval Surface Weapons Center White Oak Silver Spring, MD 20910	1
Professor Curt Wittig University of Southern CA Dept. of Electrical Engineering University Park Los Angeles, CA 90007	1	Dr. C.S. Coffey Naval Surface Weapons Center Code R13 White Oak Silver Spring, MD 20910	1

INIT

6/81

DISTRIBUTION LIST

	<u>No. Copies</u>
Dr. W. L. Elban Code R13 Naval Surface Weapons Center White Oak Silver Spring, MD 20910	1
Mr. K.J. Graham Naval Weapons Center Code 3835 China Lake, CA 93555	1
Dr. B. Junker Office of Naval Research Code 421 Arlington, VA 22217	1
Prof. H.A. Rabitz Department of Chemistry Princeton University Princeton, NH 08540	1
Dr. M. Farber Space Sciences, Inc. 135 West Maple Avenue Monrovia, CA 91016	1
Mr. M. Hill SRI International 333 Ravenswood Avenue Menlo Park, CA 94025	1
U.S. Army Research Office Engineering Division Box 12211 Research Triangle Park, NC 27709	1
U.S. Army Research Office Metallurgy & Materials Sci. Div. Box 12211 Research Triangle Park, NC 27709	1
Professor G.D. Duvall Washington State University Department of Physics Pullman, WA 99163	1

DATE
ILME